

SIGNAL & DATA ANALYSIS IN NEUROSCIENCE 2017 POISSON POINT PROCESS

Ayala Matzner

biu.sigproc@gmail.com

Poisson point process

- The probability of an event to happen is independent of previous events.
- It is totally dependent on the p.d.f that govern the process.
- $P(t_n | t_1, \dots, t_{n-1}) = P(t_n)$
- Homogeneous Poisson process: Constant firing rate r .
- Inhomogeneous Poisson process: Time dependent firing rate $r(t)$.

3

Homogeneous Poisson process

- The **mean** firing rate of the neuron is constant $r(t) = r$
- Every sequence of n spikes over a fixed time interval T has an equal probability. But not equal probability across different n 's !
- We call this probability $P_T[n]$ – the probability of getting n spikes in time span T .

$$P_T[n] = \frac{(rT)^n}{n!} e^{-rT}$$


Poisson distribution

$$\lambda = r \cdot T \quad P(n) = \frac{\lambda^n}{n!} e^{-\lambda}$$

• λ is the average events n in the interval T

• Mean = Variance = $rT = \lambda$

MATLAB - `poisspdf(a, lambda)`
`poisspdf(1:20, 10)`



Terms: definitions and properties

Term	Definition
Inter-spike interval (ISI)	<p><u>Homogeneous Poisson process:</u></p> <p>Discrete: $P(\text{ISI} = k \text{ intervals}) = (1-p)^{k-1} \cdot p$</p> <p>Continuous: $P(N(t, t + \tau) = k) = \frac{e^{-\lambda\tau} (\lambda\tau)^k}{k!}$ $P(\text{ISI} = \tau) = P(N(0, \tau - dt) = 0) \cdot P(N(\tau - dt, \tau) = 1)$</p>
Coefficient of variation (Cv)	$C_v = \frac{\text{std}(\text{ISI})}{E(\text{ISI})}$
Fano factor (FF)	$\text{FF} = \frac{\text{var}(\text{spike count over } T)}{E(\text{spike count over } T)}$

Properties of Poisson distribution

• For a homogeneous Poisson process

$$\mu = \sigma^2 = rT = \lambda$$

• For a homogeneous Poisson process

$$\text{FF} = C_v = 1.$$


FF=1 & Cv=1 is a necessary, though not sufficient, condition to identify a Poisson spike train.

• For a regular process FF = Cv = ?

• Cv > 1 → higher order process.

Example: Exam 2007

- The ISI histogram of a neuro ganglion is plotted below:



- Calculate the CV, compare to Poisson.
- Estimate the Fano factor (100msec window for rate calculation), compare to Poisson.

Solution

- Let us derive **Cv**:
 $E[ISI] = 0.5 \cdot 9 + 0.5 \cdot 10 = 9.5$
 $var[ISI] = 0.5(9 - 9.5)^2 + 0.5(10 - 9.5)^2 = 0.5^2$

$$C_V = \frac{\sqrt{var[ISI]}}{E[ISI]} = \frac{\sqrt{0.5^2}}{9.5} < 1$$

For a Poisson process $C_V \rightarrow 1$. In our case the process is close to regular.

- Let us estimate **FF** over $T = 100\text{msec}$ windows:
 $E[\text{\#spike over } T] = T/E[ISI] = 100/9.5 \sim 10.5$
 $\min[\text{\#spike over } T] = 100/10 = 10$
 $\max[\text{\#spike over } T] = 100/9 \sim 11$
 $var[\text{\#spike over } T] \sim 0.5^2$ (We can estimate without exact values)

$$FF = \frac{var[\text{\#spikes over } T]}{E(\text{\# spikes over } T)} \approx \frac{0.5^2}{10.5} \ll 1$$

Example: Exam 2006

- Neurons in the Singelitis Non-Exactis nucleus of the Mountain Elf fires a spikes with intervals of either ($p=0.5$) 100ms or ($p=0.5$) 10ms. Calculate the Coefficient of variation of the neurons. Explain the results and their relation to a Poissonian neuron firing at the same rate.

Solution

• Cv definition: $C_v = \frac{\text{std}(\text{ISI})}{E(\text{ISI})}$

$$E(\text{ISI}) = 0.5 \cdot 100[\text{msec}] + 0.5 \cdot 10[\text{msec}] = 55[\text{msec}]$$

$$\text{var}(\text{ISI}) = 0.5 \cdot (100 - 55)^2 + 0.5 \cdot (10 - 55)^2 = 45^2$$

$$C_v = \frac{45}{55} = 0.81$$

- Note that Cv is high although the neuron is not Poisson.

Example: 2005 demo exam

- A pacemaker neuron fires 10 spikes/s in a fairly regular manner ISI=N(100ms,5ms) draw the ISI distribution and estimate the Fano factor and calculate the coefficient of variation.

Normal distribution: $f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}}$

Solution

• Normal distribution: $f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}}$

• $C_v = \frac{\text{std}(\text{ISI})}{E(\text{ISI})} = \frac{5}{100}$

- Estimating **FF** over T = 100msec windows:
 E[#spikes]: p(0 spikes) ~ p(1 spike) ~ 0.5
 Assuming 2 STDs (10 ms):
 min [#spike over T] = 100/110 ~ 0.
 max [#spike over T] = 100/90 ~ 1.
 var[#spike over T] ~ 0.5^2
 FF = 0.5

Generation of a Poisson process

- Common way: Translate the Poisson RV into a binomial RV in each bin, Simulate the probability for a spike using uniform distribution (MATLAB rand function).
- Using exponential distribution to evaluate ISI.
